

THE USE OF HYDROGEN PEROXIDE AND SILVER NANOPARTICLES IN HORTICULTURE

Leszek ORLIKOWSKI^{**}, Lidia SAS-PASZT*, Adam WOJDYŁA, Teresa ORLIKOWSKA^{**}

^{**}Professor emeritus

National Institute of Horticulture, Poland

Received: August 2023; Accepted: December 2023

ABSTRACT

Both hydrogen peroxide and silver can oxidize organic and inorganic molecules, and this causes them to affect the metabolism of living organisms in many ways. The paper presents examples of the impact of H₂O₂ and silver on stimulating plant growth and development and increasing plants' resistance to biotic and abiotic stresses. The most underlined proposal for application in horticulture is the control of microorganisms during cultivating and storing vegetables, fruits, and flowers, aiming to replace synthetic pesticides. Preparations containing H₂O₂, silver, or both components can be widely used in horticulture for plant protection, in the form of spraying and soaking seedlings, to protect them during the time of storage, for disinfecting tubers, bulbs, and rhizomes before planting, for fogging potatoes and root vegetables during storage, for quick healing of wounds on the roots and aboveground parts of plants, after cutting and in the case of frost damage and injuries caused by winds, for disinfecting seeds, and, as stimulants of plant development and inducers of resistance to biotic and abiotic stresses. However, their practical use depends on obtaining the legislator's consent for their broader use in horticultural production.

Key words: hydrogen peroxide, silver nanoparticles, plant protection, control of microorganisms, plant stimulants, resistance induction

INTRODUCTION

The consistently increasing concern for the quality of plant products and the production environment in agriculture and horticulture has led to the withdrawal of many pesticides from cultivation. Unconventional plant protection products that do not pollute the produce nor the environment or pollute to a much lesser extent are being sought. Among them, some hopes are associated with hydrogen peroxide stabilized with silver, known in Poland as Huwa San TR-50 and Bisteran preparations. Huwa San TR-50, containing 50% hydrogen peroxide and 0.036% colloid silver, was approved for use in Poland by the Minister of Health No. 4236/10 of 2010 as a product for surface disinfection and by the decision of the Minister of Agriculture and Rural Development No. S-253e/17 of 2017 as a plant growth stimulant and for quick healing of plant wounds. Both components, hydrogen peroxide and silver, can negatively affect

microorganisms. However, when combined in a formulation, silver is most often considered a factor stabilizing hydrogen peroxide, slowing down its disintegration, thus extending the durability of the preparation. The mechanism of their impact will be discussed separately below.

CHARACTERISTICS OF HYDROGEN PEROXIDE AND ITS INCREASING IMPORTANCE IN THE GLOBAL ECONOMY

Hydrogen peroxide (H₂O₂), identified in 1818 by Louis Thénard (after Linley et al. 2012), is an inorganic chemical substance whose molecule differs from water by an additional oxygen atom. Oxygen is quickly released, oxidizing organic and inorganic molecules and participating in many metabolic changes in the cells of living organisms. H₂O₂ is easily soluble in water. At an average room temperature,

*Corresponding author
e-mail: lidia.sas@inhort.pl

it forms a thick, colorless liquid, gradually changing color to blue with increasing concentration. Hydrogen peroxide is more stable in acidic and neutral environments than in alkaline. Thanks to its properties, H_2O_2 has several applications in medicine, cosmetics, food, chemistry, paper, and textile industries and is commonly used to disinfect water (Linley et al. 2012). It is considered a “green industrial oxidant” because it leaves no harmful residues and quickly breaks into water and oxygen molecules. The properties of H_2O_2 have resulted in a growing demand for this compound, and its consumption has significantly increased in the last three decades (Ciriminna et al. 2016). Hydrogen peroxide is ubiquitous in the Earth’s atmosphere, human and animal organisms, and plants. In plants, it is produced as a byproduct of photosynthesis, photorespiration, fatty acid oxidation, and electron transport in mitochondria (Kuźniak & Urbanek 2000). The interest in using this compound in agriculture is related to its direct effects in controlling harmful microorganisms, modifying the internal resistance of plants to biotic and abiotic stresses, and impacting plant growth. The oxidizing effect of hydrogen peroxide is widely used in the disinfection of water, municipal and industrial wastewater, and in medicine to disinfect damage to the skin, surfaces, rooms (usually in volatile form), and medical equipment, replacing chlorine compounds and formalin. In plant production, H_2O_2 is proposed as a compound to control viral, bacterial, and fungal pathogens and plant parasitic pests, disinfect storage rooms, and protect plant products destined for storage (McDonnell 2014; Anonymous 2015). The liquid form affects easily accessible surfaces, while the gaseous form allows penetration into hard-to-reach places, e.g., in storage rooms.

The constant presence of hydrogen peroxide in the tissues of plants and animals prompted biologists to explain its role in the life of plants and microorganisms, which resulted in many experiments. For example, its participation in plant resistance to pathogens and response to mechanical damage and sudden temperature drops was studied by Juven and Pierson (1996), Orozco-Cárdenas et al. (2001), Baker and Orlandi (1995), and many others.

MECHANISM OF ACTION OF HYDROGEN PEROXIDE AND ITS EFFECT ON MICROORGANISMS AND PLANTS

Hydrogen peroxide plays a vital role in the life of plants because it is an essential participant in metabolism and processes related to plants’ response to biotic and abiotic stresses. As a result of the responses associated with the activation of immune functions, there are, among others, reinforced cell walls, enhanced plant growth and development, production of growth regulators, and phytoalexins (Dat et al. 2000; Kuźniak & Urbanek 2000; Quan et al. 2008). In these responses, H_2O_2 is a transmitter of signals initiating or controlling metabolic processes (Cheeseman 2007; Ślesak et al. 2007; Khan et al. 2018). The essence of the molecule is its high reactivity in the cell environment, which results in the formation of reactive oxygen and hydroxyl groups, which are strong oxidants (Linley et al. 2012). The high reactivity of OH radicals in the biological environment causes a cascade formation of other reactive oxygen species (ROS), affecting the chemical transformations of several organic compounds; as a result, the role of H_2O_2 in plant life involves many processes in which it can be formed or broken down, being both a substrate and a signal carrier (Cheeseman 2007; Quan et al. 2008; Petrov & Van Breusegem 2012; Khan et al. 2018). These reactive oxygen compounds can be cytotoxic and genotoxic (Abdollahi & Hosseini 2014; Ahire & Mishra 2022). Too high concentration of reactive oxygen (ROS) leads to oxidative stress that harms plants. Plants have mechanisms to annul excess H_2O_2 . Plant cells protect themselves against hydrogen peroxide through catalase activity and, in the event of DNA damage, through its rapid repair. The small dimensions of H_2O_2 particles enable easy penetration through the cell membranes of microorganisms, showing a biocidal effect against a broad spectrum of bacteria, fungi, and viruses, but also against harmful protein particles, such as prions (Żegliński 2006). At a concentration of 3%, this compound is known as hydrogen peroxide, which destroys naked cells, while at higher concentrations of 10–30%, called perhydrol, it also kills bacterial and fungal spores.

The biocidal mechanism consists in the fact that H_2O_2 penetrates cells and is decomposed thereby by protective enzymes, e.g., catalase or reducers, such as metal ions, causing the release of hydroxyl radicals affecting life-critical compounds. The main biocidal effect is lipid membranes' destruction and DNA strands' breaking. Imlay and Linn (1988) believe that H_2O_2 causes DNA damage at lower concentrations, but at higher concentrations, it also damages other cell structures, and the degree of injury is a function of exposure time. Brandi et al. (1991) showed that lower concentrations of this compound caused morphological changes in bacterial cells (e.g., filamentation) and higher cell shrinkage, which may result from damage to internal structures.

Considering the small number of agents approved for plant protection in organic farming and the complete harmlessness of hydrogen peroxide to humans and the environment because there are no harmful residues, only oxygen and water, preparations based on this compound have been introduced for practical use.

EXAMPLES OF THE USE OF HYDROGEN PEROXIDE IN HORTICULTURE

For the decontamination of water, soil, tools, and packaging

Using this compound in gaseous form is more effective than in liquid form, as it disinfects faster at low and high temperatures and, therefore, can be applied in lower concentrations than when administered in liquid form (McDonnell 2014). According to the cited author, using H_2O_2 does not risk harmful microorganisms becoming resistant to it because its action is nonspecific, unlike that of antibiotics. According to Nikkhah et al. (2010), hydrogen peroxide is one of the most commonly used disinfectants for fruit packaging. Linley et al. (2012) indicate the possibility of using the agent for disinfecting tools and water, including drinking water, and in the processing of agricultural products, for disinfecting rooms, greenhouses, plastic tunnels, and storage rooms. As one of the significant advantages of the agent, the authors indicate the safety of its use, the lack of unpleasant odor, and the possibility of using the disinfected rooms

soon after treatment. In a concentration from 3 to 6% and the fogging time from 10 minutes to 10 hours, depending on the intensity of contamination and the variety of microorganisms, H_2O_2 can be used to disinfect garden substrates, seedlings, and the root system of fruit and ornamental shrubs and trees.

When decontaminating soil and reusing growth substrates, H_2O_2 decomposes pesticide and herbicide residues by oxidizing them (Newman 2004). Its effectiveness is limited in a high organic substrate. ZeroTol is recommended as a pesticide for use in organic crops and as a sterilizer for the disinfection of greenhouses and gardening equipment to protect plants against bacteria, fungi, and viruses (Miyasaki et al. 1986). Rapid wound healing has been observed using H_2O_2 as a fogging agent on plant material with new lesions. Vänninen and Koskula (1998) showed that H_2O_2 could be used as a spray on seedlings and older cucumber plants grown under mineral wool covers as an algicide against green algae, on which water moths (*Scatella stagnalis*) feed. Reciclean containing H_2O_2 reduced algae growth by 40–60%, thus reducing the number of water moths by 73–92%. The agent also minimized algae growth in the water and on the surface of other substrates, primarily when used in the initial stage of algal development.

According to Japhet et al. (2022), the constant addition of H_2O_2 to water in a closed circuit allows it to water plants, inhibiting clogging and biofilm formation in the irrigation system. Barta and Henderson (2000) proposed use of H_2O_2 as an alternative to conventional bleach to disinfect hydroponic plant growth systems. In the concentration 0.5% reduced microbial population to near zero but it should be repeated during cultivation. Bosmans et al. (2016) showed that H_2O_2 can effectively reduce biofilm formation by rhizogenic bacteria, but its concentration should be adapted to the particular *Agrobacterium rhizogenes* strain. Parke and Fisher (2012), researching the possibilities to propagate plants with disinfecting water, indicate that, apart from chlorine and copper compounds, hydrogen peroxide is particularly effective when the *Pythium* and *Phytophthora* genera are present in the production environment. Stewart-Wade (2011) suggests the elimination of organic residues from the water used in plant production on which microorganisms

live, which makes it possible to almost wholly avoid the development of potential pathogens. Fredrickson (2005) indicated that adding hydrogen peroxide to water used for watering plants removed chlorine and sulfur residues, components of pesticides, and fertilizers from the soil, and the released oxygen caused faster decomposition of dying roots, eliminating potential plant pathogens and algae growth. Consequently, such action of the agent affected the increase in yield and plant quality. The agent is also recommended for treating seeds, soaking or watering seedlings, bulbs, and tubers, and spraying crops in the field and under covers. In a study by El-Mougy et al. (2008), hydrogen peroxide was a helpful agent for postharvest protection of strawberry fruits against *Botrytis cinerea* and disinfection of water, surfaces, and tools.

In the protection of plants against diseases and pests

The first pesticides containing hydrogen peroxide were registered in 1977 (Anonymous 2015). According to the United States Environmental Protection Agency (EPA 2014) and the Canadian General Standards Board (CGSB 2011), by 2014, there were 167 hydrogen peroxide-based agents registered in the US. Currently, many scientific reports prove the effectiveness of H₂O₂ as a biocide.

The protection of plants against diseases and pests as a result of H₂O₂ application is associated not only with the direct attack and destruction of microorganisms by this compound but also with its impact on plant metabolism, which causes the plant to acquire defensive capabilities as a result of hypersensitivity responses and systemic acquired resistance (SAR). Applying H₂O₂ mimics the process generated by plants in response to pathogen attack by rapidly producing reactive oxygen, known as the “oxygen burst” (Lei et al. 2014). The cited authors observed that *Arabidopsis* cells attacked by aphids accumulated H₂O₂, leading to a hypersensitivity reaction and the cells’ death. Numerous papers, including Bestwick et al. (1997), showed the appearance of H₂O₂ at the sites of penetration by bacterial pathogens. Small H₂O₂ molecules, after penetrating cell membranes, quickly decompose into reactive oxygen and water upon contact with biological material and show a biocidal effect on a broad spectrum of bacteria, fungi,

and viruses. Baatout et al. (2006) report that H₂O₂ increases the permeability of cell membranes, increasing with the concentration of the compound, and Qin et al. (2011) emphasize its negative impact on the mitochondria of fungal proteins.

In the scientific literature, it is possible to find both reports on experiments with H₂O₂ and recommendations for using it to fight pathogenic microorganisms and parasites. For example, hydrogen peroxide was found effective against several fungal pathogens, such as downy mildew (*Pseudoperonospora cubensis*) on cucurbits (Kuepper 2003), potato late blight (*Phytophthora infestans*) (Kuepper & Sullivan 2004), and apple scab (*Venturia inaequalis*) (Phillips 2005). Hydrogen peroxide has been recommended in organic tomato production (Diver et al. 1995). In addition, H₂O₂ was effective in controlling tomato wilt (*Ralstonia solanacearum*) (Hong et al. 2013), potato blackleg (*Dickeya solani*), pepper golden mosaic (PepGMV) (Mejía-Teniente et al. 2019), European stone fruit yellows (Musetti et al. 2005), the nematodes *Meloidogyne javaica* (Karajeh 2008); Jansen et al. 2002; Bolm et al. 2004), *Bactrocera* spp. flies on *Syzygium samarangense* (Khandaker et al. 2018). Hafez et al. (2020) reported that in an open-field experiment, H₂O₂ limited the infection of pear trees with *Erwinia amylovora*. Thakulla et al. (2022) studied the addition of hydrogen peroxide in the form of ZeroTol preparations (H₂O₂ + peracetic acid) and PERpose Plus (H₂O₂ + hydrogen dioxide) in hydroponic tomato cultivation. They found adverse effects of both preparations on the growth, dry and fresh weight of algae but a positive impact on tomato plants. The US organic farming guidelines allow the use of hydrogen peroxide-based disinfectants, fungicides, and algicides (EPA 2009).

Despite many reports on the positive effect of H₂O₂ on reducing plant diseases, there are doubts about whether this compound can be recommended for large-scale production. Miller (2006) did not recommend H₂O₂ as a biopesticide due to its short-term activity and assumed that this compound could be effectively used only for surface disinfection and water decontamination. However, its use against plant pathogens is risky, especially in field production.

As an inducer of plant resistance to pathogens and abiotic stress factors, in the stimulation of plant growth and development

H₂O₂ participates in plant metabolism in many ways, including activating plant genes related to resistance (Orozco-Cárdenas et al. 2001). An increased concentration of this compound was found in places of infection and damage, where it acts as a direct biocide and a factor accelerating wound healing (Orozco-Cárdenas et al. 2001). This process was observed within 4 hours of leaf tissue damage (Olson & Verner 1993). In addition, more active oxygen in the tissues stimulates metabolism and, depending on the conditions, acts as a transmitter of signals initiating various biochemical reactions. In this role, H₂O₂ activates the processes leading to damage regeneration, stimulating growth, and making plant organisms resistant to biotic and abiotic stresses. These activities are initiated by releasing oxygen, similar to an oxygen burst, which manifests the initiation of plant defense reactions in response to biotic and abiotic stress factors. For example, an increased oxygen concentration was observed in tomatoes at the inoculation site with the fungus *Cladosporium fulvum* (Borden & Higgins 2002). Chen et al. (1993) believe that salicylic acid, as an inducer of plant resistance to adverse environmental factors, limits the activity of catalase, which leads to the accumulation of hydrogen peroxide, thereby enhancing its effect. The consequence is the thickening of cell walls and the activation of genes that can induce systemic resistance of plants to pathogens. For example, H₂O₂ induced the resistance of damaged fruits to infection by *Penicillium* spp. (Janisiewicz et al. 2016). Increased amounts of H₂O₂ accumulated in places of mechanical damage to apples and citrus fruits resulted in injury healing (Macarisin et al. 2011). One of the factors inducing resistance is the stimulation of lignin synthesis, which is deposited in the cell walls of infected plants, preventing further pathogen spread (Ros Barceló 1998; Hückelhoven et al. 1999; Płazek 2011). A similar response was found by Martinez et al. (2000) in the case of cotton infection by *Xanthomonas campestris* pv. *malvacearum*. H₂O₂ stimulates wound healing (Ros Barceló 1998). External application of H₂O₂ in amounts not causing oxidative stress eliminated the

toxic effects of stress caused by copper, nickel, salinity, drought, and low and high temperatures, among others. Uchida et al. (2006) demonstrated the biosynthesis of osmoprotectants in plants treated with H₂O₂ in response to abiotic stresses. Szpunar-Krok et al. (2020) reported a positive effect on the efficiency of the potato photosynthetic apparatus using H₂O₂ at a concentration of 1%. Adding H₂O₂ to the soil increased photosynthetic activity and, in the end, the amount of soluble sugars in melon fruit (Ozaki et al. 2009).

Torres et al. (2003), researching the resistance of apples to *Penicillium expansum*, which causes fruit rot during storage, found an increase in the concentration of hydrogen peroxide and reactive oxygen species (ROS) in the harvested fruits as a result of artificial inoculation with the pathogen, followed by an increase in the resistance to it. Bestwick et al. (1997) found a hypersensitive response in lettuce to *Pseudomonas syringae* pv. *phaseolicola* as early as 5–6 hours after cell inoculation with the bacterium. Olson and Verner (1993) and Wu et al. (1997) showed that the agent could make potato plants resistant to several species of pathogens, causing an even several-fold increase in the concentration of salicylic acid in the affected tissues and an increase in the amount of lignin in the shoots and roots. Research by Türküsay and Tosun (2005) showed that using hydrogen peroxide for spraying tomato plants infected with *Clavibacter michiganensis* ssp. *michiganensis* can induce resistance to this bacterium. In a study conducted on cucumbers, Lee et al. (2004) showed that when there was a sharp drop in temperature, hydrogen peroxide accumulated in the cytoplasmic membranes of root cells, and the water uptake by the root system decreased by half. This action of the agent affects limiting the freezing of plants. According to Mejía-Teniente et al. (2019), exogenous application of H₂O₂ induced pepper plant resistance to PepGMV. Borden and Higgins (2002) showed that reactive oxygen species formed in tomatoes infected with *Cladosporium fulvum* strongly limited plant colonization by this pathogen. Price and Lee (1970) found that the bacterium *Lactobacillus plantarum* produced hydrogen peroxide, which strongly inhibited the growth of *Pseudomonas* spp. and other pathogenic bacteria.

Research by Webber et al. (2007) showed that foliar and soil applications of H₂O₂ accelerate the development of the root system and increase the number of nasturtium flowers. The authors believe this is related to eliminating soil pathogens, although more oxygen can accelerate faster root growth. Experiments conducted by Khandaker et al. (2018) showed a positive effect of H₂O₂ on the growth and development of wax apple (*Syzygium samarangense*). Spraying trees with hydrogen peroxide in concentrations from 0.05% to 0.2% at weekly intervals increased the number and size of fruit on the trees, reduced fruit drop, and increased fruit yield and quality. The fruits had a higher content of potassium, anthocyanins, and carotene, on average about 60%, but also a higher amount of flavonoids, sugar, proteins, and antioxidants. In the experiments by López-Delgado et al. (2005), spraying potatoes with hydrogen peroxide from the 3rd to the 10th week of their cultivation increased the thickening of shoots. It caused a significant increase in lignin in the conductive bundles and faster growth of plants. Al-Mughrabi (2010) showed that fogging potato tubers during storage and treating them with H₂O₂ before planting resulted in better germination, increased shoot number, and a higher yield of better quality tubers. On the other hand, Orozco-Cardenas and Ryan (1999) emphasize that an agent present in the conductive bundles can inhibit the movement of harmful viruses, bacteria, and fungi. It is also essential to use H₂O₂ to quickly heal potato tubers or other plants damaged during harvest and transport (Orozco-Cárdenas et al. 2001). In the technology developed by Afek et al. (2000) and Al-Mughrabi (2007), the agent was used to extend potato storage for up to 6 months.

In improving the storage

Hydrogen peroxide has also found valuable applications in spraying, bathing, or gasification in storing vegetables, fruits, and flowers. It is used to remove bacterial and fungal contamination from the surface of fruits and vegetables and as an addition to the solution in which flowers are stored. In the storage of citrus fruit, it was used as a disinfectant, replacing the previously used formaldehyde (Smilanick et al. 2014). Bathing apples in an aqueous solution of 1% H₂O₂ reduced the *Escherichia coli* population (Sapers & Sites 2003).

However, the same treatment was ineffective for decontaminating melons, meaning that the concentration and duration of exposure must depend on the fruit surface area and the plant growing conditions (Ukuku 2004). On the other hand, bathing whole melon fruit in 2.5% and 5% solutions effectively removed *Salmonella* spp. from the surface, while in the storage of cut fruit, the fruit had to be bathed whole and then cut. Using H₂O₂ at 1% and 1.5% concentrations on harvested strawberry fruit protected them with a 90% effectiveness against *Botrytis cinerea* and *Rhizopus stolonifer* (El-Mougy et al. 2008). However, hydrogen peroxide can have adverse effects, e.g., browning that reduces the quality of fruit, browning of lettuce leaves, and degradation of anthocyanins in berries (Nikkhah et al. 2010; Pietrysiak et al. 2019). H₂O₂ applied in gaseous form minimized the infection of grapes by *Botrytis cinerea* (Forney et al. 1991). Washing mushrooms in a 5% H₂O₂ solution immediately after harvesting allowed them to be stored at 4°C without reducing their quality (Sapers et al. 2006). Applying H₂O₂ protected peach flesh against darkening, gradually reducing it (Ding et al. 2009). When stored for four weeks, pepper fruits bathed in a 15% hydrogen peroxide solution maintained high fruit quality, reduced weight loss, and minimized putrefaction (Bayoumi 2008).

In potato storage, it is crucial to maintain their good health by minimizing the development of possible pathogens transferred onto the tubers and preventing the growth of sprouts (Afek et al. 2000; Kleinkopf et al. 2003). Research by Afek et al. (2000, 2001) and Al-Mughrabi (2007, 2010) indicates the possibility of using hydrogen peroxide (HPP with an unknown mixture of stabilizers, and StorOx with 27% H₂O₂) to minimize the development of pathogens causing wet rot of tubers (*Erwinia carotovora*), alternariosis (*Alternaria solani*), fusariosis (*Fusarium sambucinum*), powdery scab (*Helminthosporium solani*), dry rot (*Phytophthora erythroseptica*), late blight (*P. infestans*), and *Pythium ultimum*. Gachango et al. (2012) confirmed the effectiveness of H₂O₂ in protecting potatoes against *F. sambucinum*, *P. erythroseptica*, *P. infestans*, and *P. ultimum* in storage. They showed that H₂O₂ completely inhibited the development of *P. infestans*, with an effectiveness of four times greater

for *F. sambucinum* and six times greater for *P. ultimum* on tubers, compared to the control. Czajkowski et al. (2013) indicated that among the several agents tested for potato protection against blackleg during storage, only H₂O₂ did not cause toxicity to plants.

MECHANISM OF ACTION OF SILVER AND ITS EFFECT ON MICROORGANISMS

Silver has accompanied man since the dawn of civilization. First, it was used to make jewelry, the contact with which was supposed to guarantee health and vigor (Lara et al. 2011). In ancient Hindu medicine, silver was described as a therapeutic agent in the fight against many diseases. In modern medicine, there is a growing interest in silver as a helpful factor in treating bacterial infections of various origins, especially in the fight against multidrug-resistant bacteria. Silver has an antimicrobial effect both in the ionized form (given in the form of salt) and as nanoparticles (<100 nm). For this reason, it is used more and more commonly to disinfect water and as an additive to fabrics, cosmetics, plastics (Lara et al. 2011). The role of silver in the fight against microorganisms increased after it was possible to produce it in the form of nanoparticles and thus replace the ionic form. According to Durán et al. (2016) and Mufamadi and Mulaudzi (2019), silver nanoparticles can affect bacteria differently. For example, (i) as positively charged, it attaches itself to negatively charged bacterial cell walls, creates pores in them, penetrates inside, and binds to DNA bases, causing damage to their function (inhibition of replication); (ii) produces reactive oxygen (ROS) that changes the permeability of the walls and the outflow of intracellular fluids, leading to cell destruction, causing disruption in respiration, binding to the chromosome and causing its damage; (iii) binds to ribosomes inhibiting protein synthesis. According to Yamanaka et al. (2005), silver ions interact with ribosomes and suppress the expression of enzymes and proteins essential to ATP production, resulting in ATP deficiency. The biocidal effect of nanosilver on viruses is based on the inhibition of nucleic acid replication (Gupta et al. 2018). The hyphae of sclerotia-forming fungi (*Sclerotinia sclerotiorum*, *Rhizoctonia solani*, and *Sclerotinia minor*) were destroyed by silver nanoparticles (AgNPs), which is manifested

as the separation of layers of the hyphal wall and then its complete disintegration, what prevented germination (Min et al. 2009). Thus, silver has a multilateral nonspecific inhibitory effect on microorganisms, interfering in a broad spectrum of biological processes (Pal et al. 2007; Ouda 2014; Kędziora et al. 2016), e.g., it destroys the structure and function of cytoplasmic membranes and inactivates the main essential enzymes. The biocidal mechanisms of silver ions and nanoparticles are similar. However, due to its size, nanosilver has a greater capacity for direct contact with a bacterial cell. It penetrates the cell more efficiently and oxidizes there, gaining biocidal ability. The antimicrobial effectiveness of AgNPs depends on the size and shape of the nanoparticles, the sensitivity of microorganisms to silver (Lara et al. 2011), and the preservation of activity for a longer time than that provided by H₂O₂ (Kędziora et al. 2016). Because nanosilver particles accumulate on the outer membranes of cells (Sondi & Salopek-Sondi 2004; Gajbhiye et al. 2009), it is effective at low concentrations (Jo et al. 2009).

The influence of nanosilver on plants is also multilateral (Mahajan et al. 2022). Silver enters plant cells through pores in the cell walls, both in the roots and leaves, travels inside the cytoplasm, and is incorporated into the metabolism of various organelles (Yan & Chen 2019). Over longer distances, it travels through the xylem, reaching all parts of the plant (Yan & Chen 2019). Depending on the form of silver (ionic or nanoparticle), its concentration, plant growth stage, and environmental conditions, its effects on plants can be positive or negative.

The positive effects include interference in the hormonal balance of the cell, primarily in inhibiting ethylene levels, which delays cell aging (used in storage and in extending the shelf life of flowers). In addition, silver increases the number and size of pores in the roots, which leads to a more efficient water and nutrient uptake and better communication with the microbiome, which increases colonization with nodule bacteria and other beneficial immune-enhancing microorganisms. Thirdly, silver generates ROS, which enables the oxidation of essential elements and increases the antioxidant reserve and the formation of chlorophyll, thus increasing biomass and the amounts of secondary metabolites (Mahajan et al. 2022).

The phytotoxicity of silver manifests itself at the morphological and physiological levels. Silver can inhibit seed germination, chlorophyll and carotenoid synthesis, and root growth, reducing leaf area and biomass production (Yan & Chen 2019). Silver can negatively affect the level of macro- and microelements by allowing them to flow out of the cell. Disturbance of root geotropism and increased reduction of auxin and increase of cytokinin level were found. Depending on the plant and growth phase, exposure to silver deregulates signal genes, increasing and decreasing their expression. The penetration of silver into the cell area leads to various types of chromosomal aberrations in the studied biological systems, such as incorrect orientation in metaphase, spindle dysfunction, fragility, fragmentation, unequal separation to the poles, scattering and lagging of chromosomes, etc. The basis of the mechanism of phytotoxicity is the oxidative stress in cells induced by ROS, increased by the presence of ionic silver released from nanosilver particles or silver compounds. Ionic silver causes an n-fold increase in H₂O₂ production, leading to many disorders in metabolism and also in the course of photosynthesis. The importance of silver to the plant cell has been extensively studied, primarily in laboratory conditions. Its comprehensive effect, including a negative impact on plants and the environment, must lead to its use being precise as to the chemical form, concentration, duration of action, stage of plant development, the intended use of the product, and the type of environment in which it will remain.

The properties of silver nanoparticles allow them to be used for the mineralization of pesticides in water purification. Nanosilver has been used chiefly in medical applications, but many reports apply to agriculture (Manimegalai et al. 2011).

EXAMPLES OF THE USE OF SILVER IN HORTICULTURE

In the stimulation of plant growth and development

Gupta et al. (2018) examined the effect of silver nanoparticles on the growth of rice seedlings. They demonstrated the stimulatory effect expressed as promotion of seedlings length and biomass. Salachna et al. (2019) found that soaking lily bulbs in silver nanoparticle solution stimulated plant growth, accelerated flowering with more flowers, and flowered longer.

Pražak et al. (2020) studied the effects of AgNPs on bean seed germination and plant growth parameters at typical and low temperatures. At low concentrations (0.25 mg·l⁻¹), nanosilver stimulated quick and uniform seed germination in the laboratory and positively impacted plant growth and development in the field experiment. A particularly pronounced positive effect was revealed during germination at low temperatures, indicating that nanosilver may activate the tolerance mechanism to environmental stresses. Higher concentrations killed microorganisms but also negatively affected plant growth.

In the fight against pathogens

Kim et al. (2012) tested the *in vitro* sensitivity to AgNPs of several plant pathogens and found that all were sensitive. However, the result depended on the type of fungus and the concentration of the preparation, with the effect increasing with the concentration (up to 100 ppm). Silver administered as AgNO₃ or nanoparticles effectively eliminated or reduced the disease on ryegrass leaves, mainly when it was applied before inoculation with the fungus (Jo et al. 2009). Field experiment results of Lamsal et al. (2011) indicate that silver nanoparticles at a concentration of 10 to 100 ppm were more effective than classical fungicides in controlling pepper anthracnose (*Colletotrichum*), mainly when used preventively. In a field experiment, Gado et al. (2016) found that AgNPs reduced the severity of a disease on roses caused by *Phragmidium* spp. by one third. According to Shoaib et al. (2022), using nanosilver to control nematodes such as *Meloidogyne incognita*, *Pratylenchus penetrans*, and *Tylenchulus semipenetrans* is very promising. Al-Azzazy et al. (2019) reported that a low dosages of AgNP reduced the population of phytophagous mites associated with tomato plants in field cultivation.

In addition to the direct impact on microorganisms, silver molecules are involved in inducing the systemic resistance of plants to pathogens (Chen et al. 1993). For example, it was found that silver nanoparticles induced resistance of *Arabidopsis* seedlings to *Pseudomonas syringae* pv. *tomato* (Chu et al. 2012).

In the storage of fruits, vegetables and flowers

Nanosilver has been used to extend the life of cut flowers, and presumably, this is its most indisputable application. Its action is related to the inhibition of the multiplication of microorganisms in the storage

solution and in the conductive bundles, through which silver reaches the flower, where it silences the ethylene biosynthesis genes, which are of great importance for the durability of climacteric plants (Basiri et al. 2011; Naing & Kim 2020). The positive impact on the durability of cut flowers has been reported in, for example, carnation (Sedaghatthoor 2015), gladiolus (Li et al. 2017), tree peony (Zhao et al. 2018), gardenia foliage (Lin et al. 2019), snapdragon (Rabiza-Świder 2020), peony (Skutnik et al. 2020a), garden cosmos (Skutnik et al. 2020b), *Thalictrum aquilegifolium* (Poniewozik et al. 2020), lisianthus (Skutnik et al. 2021), tuberose (Paul et al. 2021), *Strelitzia* (Thakur et al. 2022), alstroemeria (Anwari et al. 2022), and rose (Ha et al. 2022). Nanosilver was mainly applied as pulse treatment at 5–20 mg·l⁻¹ for 24 h or as a permanent presence in the liquid in which flower stalks were immersed. Silver has also been recommended as an application on fresh fruits and vegetables to inhibit the reproduction of microorganisms and, as a result, to extend the shelf life (Kale et al. 2021). For example, coating asparagus with a nanosilver solution delayed weight loss, the loss of ascorbic acid and chlorophyll, prevented the growth of microorganisms and, as a result, extended the shelf life by ten days when stored at 2 °C (An et al. 2008). Similar results were obtained for pears and carrots (Fayaz et al. 2009). Motlagh et al. (2012) reported positive effects of storing barberry fruits in plastic bags with the addition of nanosilver.

In *in vitro* plant cultures

Silver in ionic form and as nanoparticles were used to improve the effect of *in vitro* plant cultures (Mahendran et al. 2019; Andújar et al. 2020; Zia et al. 2020). It displays many different functions as an antimicrobial agent and plant growth stimulant. One of the first reports on the effective use of silver (in the form of AgNO₃) concerned the elimination of the bacterial vector (*Agrobacterium tumefaciens*) from rose callus. Common antibiotics caused callus blackening and impaired shoot organogenesis. Silver nitrate inhibited the proliferation of bacteria and enabled organogenesis (Orlikowska et al. 1996). Nowadays, many authors recommend the use of nanosilver for the decontamination of initial explants, e.g., *Arabidopsis* seeds (Mahna et al. 2013), nodal segments of rose shoots (Shokri et al. 2015), valerian

stems (Abdi et al. 2008), capitulum of gerbera (Fakhrfeshani et al. 2012), *Araucaria* (Sarmast et al. 2011), lemongrass (Salisu et al. 2014), *Pennisetum* (Parzymies et al. 2019), and others. Parzymies (2021) reported that nanosilver particles reduced the contamination of explants but also decreased regeneration and slowed down the growth of *Aldrovanda vesiculosa*. By contrast, Andújar et al. (2020) demonstrated that silver nanoparticles (preparation Argovit™) reduced contamination and enhanced the growth and multiplication rate of *Psidium friedrichsthalianum*. Nanosilver or AgNO₃ was added to the medium to protect cultures before contamination and to affect regeneration, multiplication, shoot rooting, and stimulating the production of secondary metabolites (Kim et al. 2017; Mahendran et al. 2019; Mahajan et al. 2022; Tung et al. 2022). Depending on the plant species and goal of *in vitro* culture, the form and concentration of the agent must be determined experimentally. The most effective *in vitro* is its inhibitory action against ethylene, especially in ethylene-sensitive plants such as cruciferous vegetables (Chi & Pua 1989; Akasaka-Kennedy et al. 2005) and potatoes (Perl et al. 1988; Ehsanpour & Nejati 2013).

RESERVATIONS REGARDING THE USE OF SILVER

The antimicrobial effectiveness of silver-containing preparations and their increasingly common use in medicine, plant and animal production, and in water disinfection as an additive to clothing and everyday products raise concerns about its impact on human health and contamination of the natural environment (Jampilek & Král'ová 2018). The ease of penetration into cells, especially in ionic form, and the comprehensive influence on cell metabolism are not limited to harmful microorganisms but all living organisms. Unlike a multicellular organism, a single cell has little chance of surviving the “invasion” of silver. However, considering the ability of this element to accumulate on the surface of cell membranes, its harmfulness to higher organisms should also be considered. Nanosilver is the most dangerous for the liver, which receives all substances that the body considers toxic. However, the lungs, blood tissue, heart, nervous and immune systems may also be at risk (Rezvani et al. 2019).

The widespread use of nanosilver means that it should be used with caution because it finds many ways to the environment, first to the production environment and then to the soil, water, and air, where it moves without restrictions. The long-term impact of this biocidal agent on microorganisms makes it necessary to consider the possibility of immunizing them against it, even though the biocidal mechanism is nonspecific. Graves et al. (2015) reported that *Escherichia coli* may become more tolerant to silver nanoparticles in the 200th generation of these bacteria. Silver also affects plants, and its exact role in plant metabolism has not been precisely determined; therefore, research is ongoing (Bapat et al. 2022). The above considerations should be considered when using silver-based preparations, although there is now a consensus that they are much less harmful than synthetic pesticides (Jo et al. 2008).

PROPOSED USE OF PREPARATIONS CONTAINING H₂O₂ AND SILVER IN HORTICULTURE

H₂O₂ has advantages that make it predestined for application to plants to combat disease-causing agents directly, but its short-lived impact is its disadvantage. It is a contact agent; it can disinfect plant surfaces after application but soon leaves the plant vulnerable to subsequent microbial attacks (Miller 2006). Currently, preparations with hydrogen peroxide also contain a stabilizer that extends its durability. The stabilizer may be a metal; most often, it is silver. However, Martin et al. (2015) believe that the antimicrobial properties of such preparations (e.g., Huwa San TR-50) are mainly due to hydrogen peroxide, and the presence of silver ions allows for more effective action of the agent on microorganisms. Martin et al. (2015) and Alkawareek et al. (2019) believe that the silver in this compound facilitates electrostatic interaction on the surface of bacterial cells, adhering to the cell walls. On the other hand, Chu et al. (2011) think silver nanoparticles are potent bacteria inhibitors with a much broader antimicrobial activity than hydrogen peroxide. Shuval et al. (2009) reported that the combined formulation of H₂O₂ and silver is 100 times more potent and long-lasting than H₂O₂ alone.

Although both components are biocidal, combining them in one preparation resulted in a synergistic effect in combating microorganisms (Alkawareek et al. 2019).

In disinfection of surfaces and storage chambers

The preparation with these two components is proposed for disinfecting swimming pool water (Martin et al. 2015), surfaces, and equipment where other agents caused corrosion of metal parts, and for disinfecting rooms without the risk of leaving an unpleasant odor in with (Linley et al. 2012). As a result of a 24-month experiment, it was found that the H₂O₂/silver formulation was thermostable and more effective at higher temperatures, which predestines the use of the preparation for disinfecting hot water. Silver is deposited on the inner surfaces of pipes. The positively charged metal binds the negatively charged bacterial walls, destroying the structure of the membranes, which leads to the penetration of H₂O₂ and the death of bacterial cells (Shuval et al. 2009). Darzi et al. (2020) reported removing CGMMV virus particles from the surface of infected seed boxes in a 1% solution of Huwa San TR 50.

Martin et al. (2015) reported that the agent used to disinfect surfaces as well as to disinfect drinking water eliminated *Legionella* spp., *Escherichia coli*, *Cryptosporidium parvum*, *Giardia* spp., *Pseudomonas aeruginosa*, *Bacillus subtilis*, *Streptococcus aureus*, and others species, without affecting the taste and smell of the water. Wilson (2011) emphasized that Huwa San TR-50 added at the concentration of 200 ppm·m⁻³ disinfected drinking water, eliminating *Legionella* spp. within 24 hours. The authors speculate that this agent added at a concentration of 15–20 ppm can also kill other species of bacteria.

In Poland, research on the activity of Huwa San TR-50 in the disinfection of water intended for irrigation of horticultural crops was conducted by Orlikowski et al. (2017a, b), showing a reduction in the population of pathogenic fungi from the genera *Cylindrocladium*, *Fusarium*, *Phytophthora* and *Pythium*. An example can be an 87.5% reduction in *Phytophthora cinnamomi* by adding 100 ml of the agent per m³ of water.

Huwa San is also a secondary disinfectant, mainly to eliminate carcinogenic trihalomethanes formed after water chlorination. The World Health Organization allows up to 100 µg of silver per liter in drinking water as the active ingredient of Huwa San TR-50 (Martin et al. 2015).

Huwa San TR-50 at a concentration of 10% was used for surface decontamination of initial pistachio shoot explants. It was less effective than HgCl₂ in eliminating surface contaminants and less damaging to the explants (Korkmaz et al. 2022).

In plant protection

Linley et al. (2012) indicate that agents based on silver-stabilized hydrogen peroxide are more and more commonly used as biocides, especially since their environmental toxicity is lower than that of synthetic pesticides. Mahesha et al. (2021), in laboratory tests, studied the effectiveness of preparation based on colloidal silver with H₂O₂ (Alstasan Silvox) against some bacterial and fungal pathogens *in vitro* and its toxicity to young cotton plants and found that toxicity to plants occurred only at a concentration of 10,000 ppm. Linley et al. (2012) emphasize the much higher activity of the gaseous agent because it reaches the smallest nooks and crannies, surrounding the hyphae and spores of harmful microorganisms.

Introducing agents containing hydrogen peroxide and colloidal silver on the Polish market (Huwa San TR-50, Bisteran) resulted in an interest in its use in protecting horticultural plants, especially ornamental ones, against the most dangerous diseases. In studies by Wojdyła (2011, 2016a, b), the use of the agent at a concentration of 0.05% and 0.1% for repeated spraying plants of cissus, geraniums, willows, and roses protected them with an effectiveness of up to 80% against diseases such as powdery mildew (*Oidium* sp. and *Podosphaera pannosa*), rose black spot (*Diplocarpon rosae*), rust (*Puccinia pelargonii-zonalis* and *Melampsora epitea*), and its effectiveness ranged from 60 to 90%. In turn, Włodarek et al. (2016) showed a 60% effectiveness of the agent in protecting brassica vegetables against black cruciferous (*Alternaria* spp.). Research by Orlikowski (unpublished) assessed the effects of the agent on minimizing the

occurrence of soil pathogens of the genera *Fusarium*, *Pythium*, and *Phytophthora* in the substrate. Abundant spraying of the infected substrate with the agent at a concentration of 1 and 2%, followed by thorough mixing, resulted in the elimination of these pathogens in approximately 90%. The agent used for once-repeated spraying of poinsettias with the first symptoms of grey mold (*Botrytis cinerea*) reduced the development of the disease by about 90%, while prophylactic spraying of rooted chrysanthemum seedlings with this agent protected them entirely against the occurrence of grey mold. Huwa San TR-50 has also proven to combat grey mold on chrysanthemum micro-seedlings acclimatized in a greenhouse (Orlikowska, unpublished data). The use of the agent in the protection of orchard plants has also been tested. Applied at weekly intervals at a concentration of 0.1% for twice-repeated spraying of plum trees with cankers developing on the trunks resulted in wound healing in about 80% within two months (Orlikowski, unpublished data). Spraying ripening raspberries with the first symptoms of the grey mold with a 0.1% agent almost wholly stopped the further development of the disease (Orlikowski, unpublished data). Włodarek et al. (2015) showed that Huwa San TR-50 used to protect *Brassica pekinensis* against *Erwinia* and *Pseudomonas* resulted in better head quality and enabled extended storage.

Soika (unpublished data) showed that spraying roses four times with Huwa San TR 50 in concentrations of 0.05% and 0.1% resulted in a 95% reduction in the rose aphid population on the shoots. The above scientist sprayed thuja plants with a 0.05% Huwa San TR-50 and observed a 60% reduction in the number of thuja beetles. Alhewairini (2017) showed that spraying cotton with an agent at a concentration of 0.1% resulted in the elimination of aphids (*Aphis gossypii*) in at least 90%. At this concentration, however, the agent did not reduce the honey bee population (*Apis mellifera lamarckii*). Observations carried out in nurseries of fruit and ornamental trees sprayed with 0.1% Huwa San TR-50 for quick healing of wounds showed that the use of the agent resulted in a drastic reduction in the occurrence of the spider mite (Orlikowski, unpublished data).

Most reports on the effectiveness of Huwa San relate to the control of spider mites on various plants: date palm (Alhewairini & Al-Azzazy 2017), *Citrus sinensis* (Al-Azzazy & Alhewairini 2019; Alhewairini 2018), *Solanum lycopersicum* (Alhewairini & Al-Azzazy 2018), olive tree (Al-Azzazy & Alhewairini 2018; Alhewairini et al. 2020), *Vitis vinifera* (Al-Azzazy & Alhewairini 2020). As Wojdyła (2011) showed, Huwa San TR-50 limits the development of foliar pathogens *Podospaera pannosa*, *Puccinia pelargonii-zonalis*, *Diplocarpon rosae* and *Oidium*, and *Podospaera pannosa* on rose (Wojdyła 2016a) and *Melampsora epitea* on willow (Wojdyła 2016b). Huwa San TR-50 reduced tomato diseases caused by *Fusarium solani* and *Rhizoctonia solani* (Hamed et al. 2022). This preparation can control *Alternaria* spp. on Chinese cabbage (Włodarek et al. 2016). Grzegorzewska and Kowalska (2013) write about its use in effectively controlling *Sclerotinia sclerotiorum* on carrots. Huwa San 25 inhibited the bacterial disease caused by *Erwinia amylovora* (Hafez et al. 2020) on pear trees while stimulating shoot growth.

The bacterium *Agrobacterium rhizogenes* causes mad root disease, mainly on cucumbers and tomatoes grown in mineral wool (Martin et al. 2015; Bosmans et al. 2016). Depending on the bacterial frequency, losses can reach up to 30%. For the protection of plants against disease-causing bacteria, it is advised to remove diseased plants immediately after noticing the disease and disinfect communication routes, vehicle tires, footwear, and hands with Huwa San TR-50. Vanlommel et al. (2020) reported that immersing mineral wool bricks once in a bath with Huwa San TR-50 before planting tomato plants reduced the occurrence of the bacterium throughout the entire cultivation cycle. However, in greenhouse tomato cultivation in Belgium, it is recommended to use Huwa San TR-50 throughout the cultivation period, primarily to disinfect the irrigation system (Anonymous 2016).

In the stimulation of plant growth

Huwa San TR-50 stimulated tomato seed germination and seedling growth (Türküsay & Tosun 2005). Stepowska's research (unpublished) on the use of the agent in the cultivation of peppers showed that Huwa San TR-50 at a concentration of 0.05%, sprayed five times on plants grown in a polytunnel, increased the

chlorophyll content in the leaves. As a result, a significant increase in fruit yield and quality were obtained. Not without significance was the beneficial effect of the agent on the soil and the substrate.

Tests of the agent for registration purposes at the National Institute of Horticultural Research in Skierniewice (2012) showed its high effectiveness in the healing of wounds. Within three months of spraying with Huwa San, the scars on damaged trees healed to about 90%. These data indicate the necessity of using Huwa San TR-50 after pruning fruit trees and shrubs to heal injuries caused by storms, post-frost cracks, and injuries caused by physiological disorders. Its use is also helpful for alleviating damage to root vegetables, brassicas, and onions suffered during harvesting and transport. The agent also has a healing effect on plant organs damaged by pests, e.g., insect caterpillars, aphids, cup mites, and spider mites (Soika, unpublished). In the case of the spider mite inhabiting chrysanthemum leaves, the pest was found dead as early as 6 minutes after spraying the plants with Huwa San TR-50 at a concentration of 0.1% (unpublished data). Wojdyła et al. (2022) showed that immersion of hyacinth tubers before planting stimulated their growth and development.

In the storage of fruit and vegetables

Aharoni et al. (1994) showed that Sanosil-25 containing 48% H₂O₂ and 0.05% silver ions, added to melon fruit baths before storage, reduced the decay caused by *Fusarium solani* and *Alternaria alternata* when incorporated into wax-covering fruits. The same preparation applied as short dipping reduced the decay of eggplant and sweet red pepper fruits caused by *Botrytis cinerea* and *A. alternata* (Fallik et al. 1994). Huwa San TR-50 was effective against *Sclerotinia sclerotiorum* on freshly cut carrot slices (Grzegorzewska & Kowalska 2013). Observations on the harmfulness of nine compounds used to protect potatoes against *Dickeya solani* showed that only Huwa San TR-50 was not phytotoxic to the tubers and strongly limited the development of black leg when tubers were soaked for 15 minutes in the solution (Czajkowski et al. 2013). Therefore, Huwa San can be commonly used to minimize the occurrence and development of pathogens during potato storage (Al-Mughrabi 2010).

The presence of organic residues and minerals introduced into the water with fertilizers favors the development of microorganisms that form the biofilm inside the ducts, whose thickness increases gradually during cultivation. Research by Armon et al. (2000) showed that Huwa San TR-50 at a concentration of 0.003% added to water throughout the plant cultivation period strongly reduced biofilm formation.

CONCLUSIONS

1. Preparations containing H₂O₂ or silver, or both components, could be widely used in horticulture for plant protection, preventively and curatively, in the form of spraying and soaking seedlings before or after digging them up, to protect them during the time of storage, for disinfecting tubers, bulbs, and rhizomes before planting, for fogging potatoes and root vegetables during their storage, for quick healing of wounds on the roots and aboveground parts of plants, after cutting and in the case of frost damage and injuries caused by violent winds, for disinfecting seeds, and, as stimulants of plant development and inducers of resistance to biotic and abiotic stresses.
2. H₂O₂ and silver can be used mainly for disinfecting water, irrigation pipes, storage rooms, greenhouses and polytunnels, pallet boxes, tools, machines, and footwear.
3. Preparations containing H₂O₂, silver, or both components could substitute for synthetic pesticides, especially in sustainable production technology. Due to the possibility of environmental contamination by silver, preparations containing this ingredient should be used cautiously and according to the manufacturer's instructions.
4. To combat microbiological contamination on initial explants *in vitro* propagation and to regulate ethylene action *in vitro*.
5. Nanoparticles of silver could be used to store cut flowers as an agent, eliminating bacteria from the storage solution, destroying internal bacteria, and reducing the concentration of ethylene, which extends flower shelf-life.

Source of funding

The research was conducted as part of a research project co-financed by the European Union under the "Cooperation" measure of the Rural Development Program 2014–2020 "Implementation of an improved product, innovative technology and production organization methods in the production of bulbous ornamental plants using high-pressure fogging of refrigerated rooms with silver-stabilized hydrogen peroxide". Contract no: 00005.DDD.6509.00019.2019.05.

REFERENCES

- Abdi G., Salehi H., Khosh-Khui M. 2008. Nano silver: a novel nanomaterial for removal of bacterial contaminants in valerian (*Valeriana officinalis* L.) tissue culture. *Acta Physiologica Plantarum* 30: 709–714. DOI: 10.1007/s11738-008-0169-z.
- Abdollahi M., Hosseini A. 2014. Hydrogen peroxide. In: Wexler P. (Ed.), *Encyclopedia of Toxicology*. Elsevier, pp. 967–970. DOI: 10.1016/b978-0-12-386454-3.00736-3.
- Afek U., Orenstein J., Kim J.J. 2001. Control of silver scurf diseases in stored potato by using hydrogen peroxide plus (HPP). *Crop Protection* 20: 69–71. DOI: 10.1016/50261-2194(00)00054-5.
- Afek U., Orenstein J., Nuriel E. 2000. Using HPP (hydrogen peroxide plus) to inhibit potato sprouting during storage. *American Journal of Potato Research* 77: 63–65. DOI: 10.1007/bf02853663.
- Aharoni Y., Copel A., Fallik E. 1994. The use of hydrogen peroxide to control postharvest decay on 'Galia' melons. *Annals of Applied Biology* 125(1): 189–193. DOI: 10.1111/j.1744-7348.1994.tb04960.x.
- Ahire V., Mishra K.P. 2022. A review of reactive oxygen species and the antioxidant system. *Cancer Medicine Journal* 5: 102–107. www.tridhascholars.org
- Akasaka-Kennedy Y., Yoshida H., Takahata Y. 2005. Efficient plant regeneration from leaves of rapeseed (*Brassica napus* L.): the influence of AgNO₃ and genotype. *Plant Cell Reports* 24: 649–654. DOI: 10.1007/s00299-005-0010-s.
- Alhewairini S.S. 2017. Innovative approach for the use of Huwa San TR 50 in controlling cotton aphids (*Apis gossypii* Glover). *Journal of Agricultural Sciences* 9: 77–85. DOI: 10.5539/jas.v9n4p77.

- Alhewairini S.S. 2018. Efficacy comparison of Huwa-San TR50, abamectin and bifenthrin for the control of the oriental spider mite *Eutetranychus orientalis* (Klein) (Acari:Tetranychidae). *Pakistan Journal of Agricultural Sciences* 55: 1003–1007. DOI: 10.21162/pakjas/18.833.
- Alhewairini S.S., Al-Azzazy M.M. 2017. A new approach for controlling the date palm mite, *Oligonychus afrasiaticus* (McGregor) (Acari:Tetranychidae) using Huwa-San TR50. *Journal of Food, Agriculture and Environment* 15: 63–67.
- Alhewairini S.S., Al-Azzazy M.M. 2018. Innovative approach for the use of Huwa-San TR50 in controlling two spotted spider mite (*Tetranychus urticae* Koch) (Acari:Tetranychidae). *Pakistan Journal of Zoology* 50: 241–247. DOI: 10.17582/journal.pjz/2018.50.1.241.247.
- Alhewairini S.S., Al-Azzazy M.M. 2020. Eco-friendly approach for the control of mites associated with olive trees in Saudi Arabia. *Pakistan Journal of Agricultural Science* 57: 1211–1217. DOI: 10.21162/PAKJAS/20.0906.
- Alkawareek M.Y., Bahloul A., Abulateefeh S.R., Alkilany A.M. 2019. Synergistic antibacterial activity of silver nanoparticles and hydrogen peroxide. *PLOS One* 14(8): e0220575, DOI: 10.1371/journal.pone.0220575.
- Al-Azzazy M.M., Alhewairini S.S. 2018. Effectiveness of Huwa-San TR50 on tomato russet mite *Aculops lycopersici* (Masse) (Acari: Eriophyidae). *Pakistan Journal of Zoology* 50(3): 869–875. DOI: 10.17582/journal.pjz/2018.50.3.869.875.
- Al-Azzazy M.M., Alhewairini S.S. 2019. Huwa-San TR50 as an acaricidal agent in controlling the citrus rust mite, *Phyllocoptruta oleivora* (Ashmead) (Acari: Eriophyidae). *International Journal of Agriculture and Biology* 22(6): 1425–1430. DOI: 10.17957/ijab/15.1217.
- Al-Azzazy M.M., Ghani S.B.A., Alhewairini S.S. 2019. Field evaluation of the efficacy of silver nanoparticles (AGNP) against mites associated with tomato plants in greenhouses. *Pakistan Journal of Agricultural Sciences* 56(1): 283–288. DOI: 10.21162/pakjas/19.8155.
- Al-Azzazy M.M., Alhewairini S.S. 2020. An alternative control strategy for grape *Erineum* mites. *Pakistan Journal of Agricultural Sciences* 57: 609–614. DOI: 10.21162/pakjas/19.844.
- Al-Mughrabi K.I. 2007. Effect of treatments of potatoes in storage and pre-planting with hydrogen peroxide (H₂O₂) on emergence and yield. *Journal of Plant Sciences* 2(6): 613–618. DOI: 10.3923/jps.2007.613.618.
- Al-Mughrabi K.I. 2010. Post harvest treatment with hydrogen peroxide suppresses silver scurf (*Helminthosporium solani*), dry rot (*Fusarium sambucinum*), and soft rot (*Erwinia carotovora* subsp. *carotovora*) of stored potatoes. *Americas Journal of Plant Science and Biotechnology* 4(Special Issue 2): 74–81.
- An J., Zhang M., Wang S., Tang J. 2008. Physical, chemical and microbiological changes in stored green asparagus spears as affected by coating of silver nanoparticles-PVP. *LWT* 41(6): 1100–1107. DOI: 10.1016/j.lwt.2007.06.019.
- Anonymous 2015. Hydrogen peroxide. Crops. Technical Evaluation Report. Pesticide Research. USDA National Organic Program, 19 p. <https://www.ams.usda.gov/sites/default/files/media/Hydrogen%20Peroxide%203%20TR%202015.pdf>
- Anonymous 2016. Polyphosphate and biocide were used to deal with contamination of the water system. Onder Glas.
- Andújar I., González N., García-Ramos J.C., Bogdanichikova N., Pestryakov A., Escalona M., Concepción O. 2020. Argovit™ silver nanoparticles reduce contamination levels and improve morphological growth in the in vitro culture of *Psidium friedrichsthalianum* (O. Berg) Nied. *SN Applied Sciences* 2; 2110; 9 p. DOI: 10.1007/s42452-020-03948-9.
- Anwari M., Hashemabadi D., Asadpour L., Kaviani B. 2022. Effect of blue light and nanosilver on vase life, antioxidant enzymes and some other physiologic parameters of *Alstroemeria* ‘Napoli’ cut flowers. *Acta Scientiarum Polonorum Hortorum Cultus*. 21: 111–122. DOI: 10.24326/asphc.2022.5.10.
- Armon R., Laot N., Lev O., Shuval H., Fattal B. 2000. Controlling biofilm formation by hydrogen peroxide and silver combined disinfectant. *Water Science Technology* 42: 187–192. DOI: 10.2166/wst.2000.0312.
- Baatout S., De Boever P., Mergeay M. 2006. Physiological changes induced in four bacterial strains following oxidative stress. *Applied Biochemistry*

- and Microbiology 42(4): 369–377. DOI: 10.1134/s0003683806040053.
- Baker C.J., Orlandi E.W. 1995. Active oxygen in plant pathogenesis. Annual Review of Phytopathology 33: 299–321. DOI: 10.1146/annurev.py.33.090195.001503.
- Bapat M.S., Singh H., Shukla S.K., Singh P.P., Vo D.-V.N., Yadav A. et al. 2022. Evaluating green silver nanoparticles as prospective biopesticides: An environmental standpoint. Chemosphere 286(Part 2): 131761. DOI: 10.1016/j.chemosphere.2021.131761.
- Barta D.J., Henderson K. 2000. Use of hydrogen peroxide to disinfect hydroponic plant growth systems. 4th Annual Conference on Life Support and Biosphere Science, Baltimore, Maryland, USA. <https://ntrs.nasa.gov/api/citations/20100039590/downloads/20100039590.pdf>
- Basiri Y., Zarei H., Mashayekhi K. 2011. Effect of nanosilver treatments on vase life of cut flowers of carnation (*Dianthus caryophyllus* cv. 'White Liberty'). Journal of Advanced Laboratory Research in Biology 2(2): 40–44. <https://journals.sospublication.co.in/ab/article/view/37>
- Bayoumi Y.A. 2008. Improvement of postharvest keeping quality of white pepper fruits (*Capsicum annuum*, L.) by hydrogen peroxide treatment under storage conditions. Acta Biologica Szege diensis 52(1): 7–15. <https://abs.bibl.u-szeged.hu/index.php/abs/article/view/2572/2564>
- Bestwick C.S., Brown I.R., Bennett M.H.R., Mansfield J.W. 1997. Localization of hydrogen peroxide accumulation during the hypersensitive reaction of lettuce cells to *Pseudomonas syringae* pv *phaseolicola*. Plant Cell 9(2): 209–221. DOI: 10.2307/3870542.
- Bolm M., Jansen W.T.M., Schnabel R., Chhatwal G.S. 2004. Hydrogen peroxide-mediated killing of *Caenorhabditis elegans*: a common feature of different streptococcal species. Infection and Immunity 72(2): 1192–1194. DOI: 10.1128/iai.72.2.1192-1194.2004.
- Borden S., Higgins V.J. 2002. Hydrogen peroxide plays a critical role in the defense response of tomato to *Cladosporium fulvum*. Physiological and Molecular Plant Pathology 61: 227–236. DOI: 10.1006/pmpp.2002.043.
- Bosmans L., Van Calenberg B., Pealeman A., Moerkens R., Wittemens L., Van Kerckhove S. et al. Efficacy of hydrogen peroxide treatment for control of hairy root disease caused by rhizogenic agrobacteria. Journal of Applied Microbiology 121: 519–527. DOI: 10.1111/jam.13187.
- Brandi G., Salvaggio L., Cattabeni F., Cantoni O., Mortelmans K.E. 1991. Cytocidal and filamentous response of *Escherichia coli* cells exposed to low concentrations of hydrogen peroxide and hydroxyl radical scavengers. Environmental and Molecular Mutagenesis 18: 22–27. DOI: 10.1002/em.2850180105.
- CGSB 2011. Organic Production Systems Permitted Substances List, 2006. Canadian General Standards Board. [retrieved 2014]
- Cheeseman J.M. 2007. Hydrogen peroxide and plant stress: a challenging relationship. Plant Stress 1: 4–15.
- Chen Z., Silva H., Klessig D.F. 1993. Active oxygen species in the induction of systemic acquired resistance by salicylic acid in plants. Science 262: 1883–1886. DOI: 10.1126/science8266079.
- Chi G.-L., Pua E.-C. 1989. Ethylene inhibitors enhanced de novo shoot regeneration from cotyledons of *Brassica campestris* ssp. *chinensis* (Chinese cabbage) in vitro. Plant Science 64(2): 243–250. DOI: 10.1016/0168-9452(89)90030-7.
- Chu H., Kim H.J., Kim J.S., Kim M.S., Yoon B.D. 2012. A nanosized Ag-silica hybrid complex prepared by gamma irradiation activities the defense response in *Arabidopsis*. Radiation Physics and Chemistry 81: 180–184.
- Ciriminna R., Albanese L., Meneguzzo F., Pagliaro M. 2016. Hydrogen peroxide: a key chemical for today's sustainable development. ChemSusChem 9: 3374–3381. DOI: 10.1002/cssc.201600895.
- Czajkowski R., de Boer W.J., van der Wof J.M. 2013. Chemical disinfectants can reduced potato blackleg caused by *Dickeya solani*. European Journal of Plant Pathology 136: 419–432. DOI: 10.1007/s10658-013-0177-8.
- Dat J., Vandenabeele S., Vranova E., van Montagu M., Inz D., van Breusegem F. 2000. Dual action of the active oxygen species during plant stress responses. Cell Molecular Life Science 57: 779–795. DOI: 10.1007/s000180050041.
- Darzi E., Lachman O., Smith E., Koren A., Klein E., Pass N., Frenkel O., Dombrovsky A. 2020. Paths of cucumber green mottle virus disease spread and disinfectant-based management. Annals of Applied Biology 177: 374–384. DOI: 10.1111/aab.12629.
- Ding Z., Tian S., Meng X, Xu Y. 2009. Hydrogen peroxide is correlated with browning in peach fruit stored at low

- temperature. *Frontiers of Chemical Engineering in China* 3: 363–374. DOI: 10.1007/s11705-009-0246-x.
- Diver S., Kuepper G., Born H. 1995. Organic tomato production. *Appropriate Technology Transfer for Rural Areas*. <https://atradev.ncat.org/wp-content/uploads/2022/06/organic-tomato-production.pdf>
- Durán N., Durán M., Bispo de Jesus M., Seabra A.B., Fávaro W.J., Nakazato G. 2016. Silver nanoparticles: A new view on mechanistic aspects on antimicrobial activity. *Nanomedicine: Nanotechnology, Biology and Medicine* 12(3): 789–799. DOI: 10.1016/j.nano.2015.11.016.
- Ehsanpour A.A., Nejati Z. 2013. Effect of nanosilver on potato plant growth and protoplast viability. *Biological Letters* 50: 35–43. DOI: 10.2478/biolet-2013-0004.
- El-Mougy N.S., El-Gamal N.G., Abdalla M.A. 2008. The use of fungicide alternatives for controlling decay of strawberry and orange fruits. *Journal of Plant Protection Research* 48: 385–395. v10045-008-0048-z.
- EPA 2009. United States Environmental Protection Agency
- EPA 2014. United States Environmental Protection Agency
- Fakhrfeshani M., Bagheri A., Sharifi A. 2012. Disinfecting effects of nano silver fluids in gerbera (*Gerbera jamesonii*) capitulum tissue culture. *Journal of Biological and Environmental Sciences* 6(17): 121–127. <https://dergipark.org.tr/tr/download/article-file/497511>
- Fallik E., Aharoni Y., Grinberg S., Copel A., Klein J.D. 1994. Postharvest hydrogen peroxide treatment inhibits decay in eggplant and sweet red pepper. *Crop Protection* 13(6): 451–454. DOI: 10.1016/0261-2194(94)90094-9.
- Fayaz A.M., Balaji K., Girilal M., Kalaichelvan P.T., Venkatesan R. 2009. Mycobased synthesis of silver nanoparticles and their incorporation into sodium alginate film for vegetable and fruit preservation. *Journal of Agriculture and Food Chemistry* 57: 6246–6252. DOI: 10.1021/jf900337h.
- Forney C.F., Rij R.E., Denis-Arrue R., Smilanick J.L. 1991. Vapor phase hydrogen peroxide inhibits postharvest decay of table grapes. *HortScience* 26(12): 1512–1514. DOI: 10.21273/hortsci.26.12.1512.
- Fredrickson B. 2005. Hydrogen peroxide and Horticulture. <https://www.quickgrow.com/hydrogen-peroxide-horticulture>
- Gachango E., Kirk W., Schafer R., Wharton P. 2012. Evaluation and comparison of biocontrol and conventional fungicides for control of postharvest potato tuber diseases. *Biological Control* 63: 115–120. DOI: 10.1016/j.biocontrol.2012.07.005.
- Gado E.A.M., El-Deeb B., Ali E.F., Mostafa N.Y., Bazaid S.A. 2016. Evaluation of silver nanoparticles for the control of *Phragmidium* species in vitro and Taif’ rose rust disease in field. *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 7(3): 886–896.
- Gajbhiye M., Kesharwani J., Ingle A., Gade A., Rai M. 2009. Fungus-mediated synthesis of silver nanoparticles and their activity against pathogenic fungi in combination with fluconazole. *Nanomedicine: Nanotechnology, Biology and Medicine* 5: 382–386. DOI: 10.1016/j.nano.2009.06.005.
- Graves J.L. Jr., Tajkarimi M., Cunningham Q., Campbell A., Nonga H., Harrison S.H., Barrick J.E. 2015. Rapid evolution of silver nanoparticle resistance in *Escherichia coli*. *Frontiers in Genetics* 6: 42; 13 p. DOI: 10.3389/fgene.2015.00042.
- Grzegorzewska M., Kowalska B. 2013. The influence of nano-silver, nano-copper and hydrogen peroxide on vegetable pathogens. *Zeszyty Naukowe Instytutu Ogrodnictwa* 21: 15–23. [in Polish with English abstract]
- Gupta N., Upadhyaya C.P., Singh A., Abd-Elsalam K.A., Prasad R. 2018. Applications of silver nanoparticles in plant protection. In: Abd-Elsalam K.A., Prasad R. (Eds.), *Nanobiotechnology Applications in Plant Protection*. Springer, pp. 247–265. DOI: 10.1007/978-3-319-91161-8_9.
- Ha S.T.T., In B.-C. 2022. Combined nano silver, alfa-aminoisobutyric acid, and 1-methylcyclopropene treatment delays the senescence of cut roses with different ethylene sensitivities. *Horticulturae* 8(6): 482. DOI: 10.3390/horticulturae8060482.
- Hafez Y.M., El-Kady E.M., Moussa Z., Elsaed N., Kotb H., Salama A.-M. et al. 2020. Exogenous application of *Bacillus subtilis* and H₂O₂ mitigated fire pear blight bacterial disease incidence in correlative with yield and fruit quality improvement. *Fresenius Environmental Bulletin* 29(7A): 6315–6327.
- Hamed K.E., Alsaif A.N., Alhewairini S.S. 2022. Soil drenching with Huwa-San TR50 for controlling *Rhizoctonia solani* (Kuhn) and *Fusarium solani* (Marti) causing damping-off disease in tomato (*Lycopersicon esculentum* (Mill)). *Pakistan Journal of Agricultural Science* 59: 319–327. DOI:

- 10.21162/pakjas/22.86.
- Hong J.K., Kang S.R., Kim Y.H., Yoon D.J., Kim D.H., Kim H.J. et al. 2013. Open access: Hydrogen peroxide and nitric oxide-mediated disease control of bacterial wilt in tomato plants. *Plant Pathology Journal* 29: 386–396. DOI: 10.5423/ppj.oa.04.2013.0043.
- Hückelhoven R., Fodor J., Preis C., Kogel K.-H. 1999. Hypersensitive cell death and papilla formation in barley attacked by the powdery mildew fungus are associated with hydrogen peroxide but not with salicylic acid accumulation. *Plant Physiology* 119(4): 1251–1260. DOI: 10.1104/pp.119.4.1251.
- Imlay J.A., Linn S. 1988. Bimodal pattern of killing of DNA-repair-defective or anoxically grown *Escherichia coli* by hydrogen peroxide. *Journal of Bacteriology* 166: 519–527. DOI: 10.1128/j.b.166.2.519-527.1986.
- Jampílek J., Král'ová K. 2018. Benefits and potential risks of nanotechnology applications in crop protection. In: Abd-Elsalam K.A., Prasad R. (Eds.), *Nanobiotechnology Applications in Plant Protection*. Springer, pp. 189–246. DOI: 10.1007/978-3-319-91161-8_8.
- Janisiewicz W.J., Nichols B., Bauchan G.R., Chao T.C., Jurick II W.M. 2016. Wound responses of wild apples suggest multiple resistance mechanism against blue mold decay. *Postharvest Biology and Technology* 117: 132–140. DOI: 10.1016/j.postharvbio.2015.12.004.
- Jansen W.T.M., Bolm M., Balling R., Chhatwal G.S., Schnabel R. 2002. Hydrogen peroxide-mediated killing of *Caenorhabditis elegans* by *Streptococcus pyogenes*. *Infection and Immunity* 70: 5202–5207. DOI: 10.1128/iai.70.9.5202-5207.2002.
- Japhet N., Tarchitzky J., Chen Y. 2022. Effectiveness of hydrogen peroxide treatments in preventing biofilm clogging in drip irrigation systems applying treated wastewater. *Journal of Bioadhesion and Biofilm Research* 38(6): 575–592. DOI: 10.1080/08927014.2022.2102903.
- Jo Y.-K., Kim B.H., Jung G. 2009. Antifungal activity of silver ions and nanoparticles on phytopathogenic fungi. *Plant Disease* 93: 1037–1043. DOI: 10.1094/pdis-93-10-1037.
- Juven B.J., Pierson M.D. 1996. Antibacterial effect of hydrogen peroxide and methods for its detection and quantitation. *Journal of Food Protection* 59(11): 1233–1241. DOI: 10.4315/0362-028x-59.11.1233.
- Kale S.K., Parishwad G.V., Husainy A.S.N., Patil A.S. 2021. Emerging agriculture applications of silver nanoparticles. *ES Food and Forestry* 3: 17–22. DOI: 10.30919/esfaf438.
- Karajeh M.R. 2008. Interaction of root-knot nematode (*Meloidogyne javanica*) and tomato as affected by hydrogen peroxide. *Journal of Plant Protection Research* 48: 181–187. DOI: 10.2478/v10045-008-0021-x.
- Kędziora A., Krzyżewska E., Dudek B., Bugła-Płoskońska G. 2016. Udział białek błony zewnętrznej we wrażliwości bakterii na nanosrebro. *Postępy Higieny Medycyny Doświadczalnej* 70: 610–617. DOI: 10.5604/17322693.1205005. [in Polish with English abstract]
- Khan T.A., Yusuf M., Fariduddin Q. 2018. Hydrogen peroxide in regulation of plant metabolism: signalling and its effect under abiotic stress. *Photosynthetica* 56: 1237–1248. DOI: 10.1007/s11099-018-0830-8.
- Khandaker M.M., Ismail S.Z., Hafiza N., Mat N., Ngah N. 2018. Effects of hydrogen peroxide and methyl eugenol on fruit growth, yield and fruit fly infestation of *Syzygium samarangense*. *International Journal of Engineering and Technology* 7(4; Special Issue 43): 54–58. DOI: 10.14419/ijet.v7i4.43.25819.
- Kim D.H., Gopal J., Sivanesan I. 2017. Nanomaterials in plant tissue culture: the disclosed and undisclosed. *Royal Society of Chemistry Advances* 7: 36492–36505. DOI: 10.1039/c7ra07025j.
- Kim S.W., Jung J.H., Lamsal K., Kim Y.S., Min J.S., Lee Y.S. 2012. Antifungal effect of silver nanoparticles (AgNPs) against various plant pathogenic fungi. *Mycobiology* 40: 53–58. DOI: 10.5941/myco.2012.40.1.053.
- Kleinkopf G.E., Oberg N.A., Olsen N.L. 2003. Sprout inhibition in storage: current status, new chemistries and natural compounds. *American Journal of Potato Research* 80: 317–327. DOI: 10.1007/bf02854316.
- Korkmaz E., Yaşar R., Soydan B., Sarpkaya K., Açar I., Aasim M. 2022. Optimization of in vitro sterilization for pistachio (*Pistacia vera* L.) rootstocks. *Anatolian Journal of Botany* 6: 1–6. DOI: 10.30616/ajb.999973.
- Kuepper G. 2003. Downy mildew control in cucurbits. *Appropriate Technology Transfer for Rural Areas*. <https://attra.ncat.org/wp-content/uploads/2022/12/downymil.pdf>
- Kuepper G., Sullivan P. 2004. Organic alternatives for late blight control in potatoes. *Appropriate Technology*

- Transfer for Rural Areas. <https://attra.ncat.org/wp-content/uploads/2022/10/lateblight.pdf>
- Kuźniak E., Urbanek H. 2000. The involvement of hydrogen peroxide in plant responses to stresses. *Acta Physiologiae Plantarum* 22(2): 195–203. DOI: 10.1007/s11738-000-0076-4.
- Lamsal K., Kim S.W., Jung J.H., Kim Y.S., Kim K.S., Lee Y.S. 2011. Application of silver nanoparticles for control of *Colletotrichum* species *in vitro* and pepper anthracnose disease in field. *Mycobiology* 39: 194–199. DOI: 10.5941/myco.2011.39.3.194.
- Lara H.H., Garza-Treviño E.N., Ixtapan-Turrent L., Singh D.K. 2011. Silver nanoparticles are broad-spectrum bactericidal and virulicidal compounds. *Journal of Nanobiotechnology* 9: 30. DOI: 10.1186/1477-3155-9-30.
- Lee S.H., Sing A.P., Chung G.Ch. 2004. Rapid accumulation of hydrogen peroxide in cucumber roots due to exposure to low temperature appears to mediate decreases in water transport. *Journal of Experimental Botany* 55: 1733–1741. DOI: 10.1093/jxb/erh189.
- Lei J., Finlayson S.A., Salzman R.A., Shan L., Zhu-Salzman K. 2014. *Botrytis* – induced kinase1 modulates *Arabidopsis* resistance to green peach aphids via phytoalexin deficient4. *Plant Physiology* 165: 1657–1670. DOI: 10.1104/pp.114.242206.
- Li H., Li H., Liu J., Luo Z., Joyce D., He S. 2017. Nano-silver treatments reduced bacterial colonization and biofilm formation at the stem-ends of cut gladiolus ‘Eerde’ spikes. *Postharvest Biology and Technology* 123: 102–111. DOI: 10.1016/j.postharvbio.2016.08.014.
- Lin S., Li H., Xian X., Lin X., Pang Z., Liu J., He S. 2019. Nano-silver pretreatment delays wilting of cut gardenia foliage by inhibiting bacterial xylem blockage. *Scientia Horticulturae* 246: 791–796. DOI: 10.1016/j.scienta.2018.11.050.
- Linley E., Denyer S.P., McDonnell G., Simons C., Mailard J-V. 2012. Use of hydrogen peroxide as a biocide: new consideration of its mechanisms of biological action. *Journal of Antimicrobial Chemotherapy* 67: 1589–1596. DOI: 10.1093/jac/dks129.
- López-Delgado H., Zavaleta-Mancera H.A., Mora-Herrera M.E., Vázquez-Rivera M., Flores-Gutiérrez F.X., Scott I.M. 2005. Hydrogen peroxide increases potato tuber and stem starch content, stem diameter, and stem lignin content. *American Journal of Potato Research* 82: 279–285. DOI: 10.1007/bf02871957.
- Macarisin D., Bauchan G., Droby S., Wisniewski M. 2011. Novel role for reactive oxygen species (ROS) in host-antagonistic yeast-pathogen interaction in post-harvest biocontrol systems. *Acta Horticulturae* 905: 113–119. DOI: 10.17660/actahortic.2011.905.11.
- Mahajan S., Kadam J., Dhawal P., Barve S., Kakodkar S. 2022. Application of silver nanoparticles in in-vitro plant growth and metabolite production: revisiting its scope and feasibility. *Plant Cell Tissue and Organ Culture* 150: 15–39. DOI: 10.1007/s11240-022-02249-w.
- Mahendran D., Geetha N., Venkatachalam P. 2019. Role of silver nitrate and silver nanoparticles on tissue culture medium and enhanced the plant growth and development. In: Kumar M., Muthusamy A., Kumar V., Bhalla-Sarin N. (Eds.), *In vitro plant breeding towards novel agronomic traits*. Springer, pp. 59–74. DOI: 10.1007/978-981-32-9824-8_4.
- Mahesha H.S., Vinay J.U., Ravikumar M.R., Visweswarasrastry S., Keerthi M.C., Halli H.M. et al. 2021. Colloidal silver hydrogen peroxide: new generation molecule for management of phytopathogens. *Horticulturae* 7: 573. DOI: 10.3390/horticulturae7120573.
- Mahna N., Vahed S.Z., Khani S. 2013. Plant *in vitro* culture goes nano: nanosilver-mediated decontamination of ex vitro explants. *Nanomedicine and Nanotechnology* 4: 2. DOI: 10.4172/2157-7439.1000161.
- Manimegalai G., Kumar S.S., Sharma C. 2011. Pesticide mineralization in water using silver nanoparticles. *International Journal of Chemical Sciences* 9(3): 1463–1471. <https://www.tsjournals.com/articles/pesticide-mineralization-in-water-using-silver-nanoparticles.pdf>
- Martin N.L., Bass P., Liss S.N. 2015. Antibacterial properties and mechanism of activity of a novel silver-stabilized hydrogen peroxide. *PLoS One* 10: 1–20. DOI: 10.1371/journal.pone0131345.
- Martinez C., Baccou J-C., Bresson E., Baissac Y., Daniel J-F., Jalloul A., Montillet J-L., Geiger J-P., Assigbetsé K., Nicole M. 2000. Salicylic acid mediated by the oxidative burst is a key molecule in local and systemic responses of cotton challenged by an avirulent race of *Xanthomonas campestris* pv *malvacearum*. *Plant Physiology* 122: 757–766. DOI: 10.1104/pp.122.3.757.
- McDonnell G. 2014. The use of hydrogen peroxide for disinfection and sterilization applications. PATAI’s

- Chemistry of Functional Groups. 1–34. DOI: 10.1002/9780470682531.pat0885.
- Mejía-Teniente L., Durán-Flores B.A., Torres-Pacheco I., González-Chavira M.M., Rivera-Bustamante R.F., Feregrino-Perez A.A. et al. 2019. Hydrogen peroxide protects pepper (*Capsicum annuum* L.) against pepper golden mosaic geminivirus (PepGMV) infections. *Physiological and Molecular Plant Pathology* 106: 23–29. DOI: 10.1016/j.pmpp.2018.11.008.
- Miller S. 2006. Can hydrogen peroxide (or dioxide-H₂O₂) control plant disease? *VegNet* 13(17). Ohio State University Extension Vegetable Crops. <http://vegnet.osu.edu>
- Min J.-S., Kim K.S., Kim S.W., Jung J.H., Lamsal K., Kim S.B. et al. 2009. Effect of colloidal silver nanoparticles on sclerotium-forming phytopathogenic fungi. *Plant Pathology Journal* 25: 376–380. DOI: 10.5423/ppj.2009.25.4.376.
- Miyasaki K.T., Genco R.J., Wilson M.E. 1986. Antimicrobial properties of hydrogen peroxide and sodium bicarbonate individually and in combination against selected oral, gram-negative, facultative bacteria *Journal of Dental Research* 65(9): 1142–1148. DOI: 10.1177/00220345860650090601.
- Motlagh N.V., Mosavian M.T.H., Mortazavi S.A. 2012. Effect of polyethylene packaging modified with silver particles on the microbial, sensory and appearance of dried barberry. *Packaging Technology and Science* 26: 39–49. DOI: 10.1002/pts.1966.
- Mufamadi M.S., Mulaudzi R.B. 2019. Green engineering of silver nanoparticles to combat plant and foodborne pathogens: potential economic impact and food quality. In: Prasad R. (Ed.), *Plant Nanobionics*. Springer, pp. 451–476. DOI: 10.1007/978-3-030-16379-2_16.
- Musetti R., Sanità di Toppi L., Martini M., Ferrini F., Loschi A., Favali M.A., Osler R. 2005. Hydrogen peroxide localization and antioxidant status in the recovery of apricot plants from European Stone Fruit Yellows. *European Journal of Plant Pathology* 112: 53–61. DOI: 10.1007/s10658-004-8233-z.
- Naing A.H., Kim C.K. 2020. Application of nano-silver particles to control the postharvest biology of cut flowers: A review. *Scientia Horticulture* 270: 109463. DOI: 10.1016/j.scienta.2020.109463.
- Newman S.E. 2004. Disinfecting irrigation water for disease management. 20th Annual Conference on Pest Management on Ornamentals. Society of American Florists, USA, 10 p. <http://www.nurserycrop-science.info/water/filtration-disinfection/other-references/newman-2004-disinfecting-irrigation-water-paper.pdf>
- Nikkhah E., Khaiamy M., Heidary R., Azar A.S. 2010. The effect of ascorbic acid and H₂O₂ treatment on the stability of anthocyanin pigments in berries. *Turkish Journal of Biology* 34(1): 47–53. DOI: 10.3906/biy-0805-14.
- Olson P.P., Verner J.E. 1993. Hydrogen peroxide and lignification. *Plant Journal* 4: 887–892. DOI: 10.1046/j.1365-313x.1993.04050887.
- Orlikowska T., Kucharska D., Nowak E., Rojek Ż. 1996. Influence of silver nitrate on regeneration and transformation of roses. *Journal of Applied Genetics* 37A: 123–125.
- Orlikowski L.B., Treder W., Ptaszek M., Trzewik A., Kowalczyk W., Łążecka U. 2017a. Necessity of disinfection water for crop irrigation. *Infrastructure and Ecology of Rural Areas* 2017(4; 1): 1387–1400. DOI: 10.14597/infraeco.2017.4.1.106.
- Orlikowski L.B., Treder W., Ptaszek M., Trzewik A., Kowalczyk W., Łążecka U. 2017b. Efektywność różnych metod odkażania w minimalizacji występowania patogenów w wodzie. 22 Sympozjum Nawadniania Roślin, Bydgoszcz-Tleń. [in Polish]
- Orozco-Cardenas M.L., Ryan C.A. 1999. Hydrogen peroxide is generated systemically in plant leaves by wounding and systemin via the octadecanoid pathway. *Proceedings of the National Academy of Sciences* 96(11): 6553–6557. DOI: 10.1073/pnas.96.11.6553.
- Orozco-Cárdenas M.L., Narváez-Vásquez J., Ryan C.A. 2001. Hydrogen peroxide acts as second messenger for the induction of defense genes in tomato plants in response to wounding, systemin, and methyl jasmonate. *Plant Cell* 13(1): 179–191. DOI: 10.1105/tpc.13.1.179.
- Ouda S.M. 2014. Antifungal activity of silver and copper nanoparticles on two plant pathogens, *Alternaria alternata* and *Botrytis cinerea*. *Research Journal of Microbiology* 9: 34–42. DOI: 10.3923/jm.2014.34.42.
- Ozaki K., Uchida A., Takabe T., Shinagawa F., Tanaka Y., Takabe T. et al. 2009. Enrichment in sugar content in melon fruits by hydrogen peroxide treatment. *Journal of Plant Physiology* 166(6): 569–578. DOI: 10.1016/j.jplph.2008.08.007.

- Pal S., Tak Y.K., Song J.M. 2007. Does the antibacterial activity of silver nanoparticles depend on the shape of nanoparticle? A study of the gram-negative bacterium *Escherichia coli*. *Applied and Environmental Microbiology* 73: 1712–1720. DOI: 10.1128/aem.02218-6.
- Parke J., Fisher P. 2012. Treating irrigation water. *Digger* 56: 41–45. Oregon State University.
- Parzymies M. 2021. Nano-silver particles reduce contaminations in tissue culture but decrease regeneration rate and slows down growth and development of *Aldrovanda vesiculosa* explants. *Applied Sciences* 11: 3653. DOI: 10.3390/app11083653.
- Parzymies M., Pudelska K., Poniewozik M. 2019. The use of nano-silver for disinfection of *Pennisetum alopecuroides* plant material for tissue culture. *Acta Scientiarum Polonorum, Hortorum Cultus* 18(3): 127–135. DOI: 10.24326/asphc.2019.3.12.
- Paul D., Jannat A., Al Mahmud A., Akhter M.J., Mahmood S. 2021. Preservative solution on vase life and quality of cut *Polianthes tuberosa* L. *Ornamental Horticulture* 27: 417–424. DOI: 10.1590/2447-536X.v27i3.2375.
- Perl A., Aviv D., Galun E. 1988. Ethylene and in vitro culture of potato: suppression of ethylene generation vastly improves protoplast yield, plating efficiency and transient expression of an alien gene. *Plant Cell Reports* 7: 403–406. DOI: 10.1007/bf00269523.
- Petrov V.D., Van Breusegem F. 2012. Hydrogen peroxide – a central hub for information flow in plant cells. *AoB Plants* 2012; pls014; 13 p. DOI: 10.1093/aobpla/pls014.
- Phillips M. 2005. *The Apple Grower: Guide for the Organic Orchardist*, 2nd ed. Chelsea Green Publishing.
- Pietrysiak E., Smith S., Ganjyal G.M. 2019. Food safety interventions to control *Listeria monocytogenes* in the fresh apple packing industry: a review. *Comprehensive Reviews in Food Science and Food Safety* 18: 1705–1726. DOI: 10.1111/1541-4337.12496.
- Płażek A. 2011. *Patofizjologia roślin*. Uniwersytet Rolniczy, Kraków, 140 p. [in Polish]
- Poniewozik M., Pogroszewska E., Rubinowska K., Dudkiewicz M., Kozak D. 2020. The effect of nanosilver on postharvest longevity of *Thalictrum aquilegifolium* L. foliage. *Acta Agrobotanica* 73(3); 7331; 11 p. DOI: 10.5586/aa.7331.
- Prażak R., Święciło A., Krzepińko A., Michałek S., Marczevska M. 2020. Impact of Ag nanoparticles on seed germination and seedling growth of green beans in normal and chill temperatures. *Agriculture* 10: 312. DOI: 10.3390/agriculture10080312.
- Price R.J., Lee J.S. 1970. Inhibition of *Pseudomonas* species by hydrogen peroxide producing Lactobacilli. *Journal of Food Protection* 33: 13–18. DOI: 10.4315/0022-2747-33.1.13
- Quan L.-J., Zhang B., Shi W.-W., Li H.-Y. 2008. Hydrogen peroxide in plants: a versatile molecule of the reactive oxygen species network. *Journal of Integrative Plant Biology* 50: 2–18. DOI: 10.1111/j.1744-7909.2007.00599.x.
- Qin G., Liu J., Cao B., Li B., Tian S. 2011. Hydrogen peroxide acts on sensitive mitochondrial proteins to induce death of a fungal pathogen revealed by proteomic analysis. *PLoS ONE*: 6(7); e21945; 14 p. DOI: 10.1371/journal.pone.0021945.
- Rabiza-Świder J., Skutnik E., Jędrzejuk A., Rochala-Wojciechowska J. 2020. Nanosilver and sucrose delay the senescence of cut snapdragon flowers. *Postharvest Biology and Technology* 165: 111165. DOI: 10.1016/j.postharvbio.2020.111165.
- Rezvani E., Rafferty A., McGuinness C., Kennedy J. 2019. Adverse effects of nanosilver on human health and the environment. *Acta Biomaterialia* 94: 145–159. DOI: 10.1016/j.actbio.2019.05.042.
- Ros Barceló A. 1998. Hydrogen peroxide production is a general property of the lignifying xylem from vascular plants. *Annals of Botany* 82(1): 97–103. DOI: 10.1006/anbo.1998.0655.
- Salachna P., Byczyńska A., Zawadzińska A., Piechocki R., Mizieleńska M. 2019. Stimulatory effect of silver nanoparticles on the growth and flowering of potted oriental lilies. *Agronomy* 9(10); 610; 14 p. DOI: 10.3390/agronomy9100610.
- Salisu I.B., Abubakar A.S., Sharma M., Pudake R.N. 2014. Study of antimicrobial activity of nano silver (NS) in tissue culture media. *International Journal of Current Research Review* 6: 13.
- Sapers G.M., Sites J.E. 2003. Efficacy of 1% hydrogen peroxide wash in decontaminating apples and cantaloupe melons. *Journal of Food Science* 68: 1793–1797. DOI: 10.1111/j.1365-2621.2003.tb12331.x.

- Sapers G.M., Miller R.L., Pilizota V., Kamp F. 2006. Shelf-life extension of fresh mushrooms (*Agaricus bisporus*) by application of hydrogen peroxide and browning inhibitors. *Journal of Food Sciences* 66: 362–366. DOI: 10.1111/j.1365-2621.tb11347.x.
- Sarmast M., Salehi H., Khosh-Khui M. 2011. Nano silver treatment is effective in reducing bacterial contaminations of *Araukaria excelsa* R. Br. var. *glauca* explants. *Acta Biologica Hungarica* 62: 477–484. DOI: 10.1556/abiol.62.2011.4.12.
- Sedaghatthoor S. 2015. Effect of wall colors and nanosilver treatment on the vase life of cut carnation “Express”. *Journal of Ornamental Plants* 5(1): 1–6. https://jornamental.rasht.iau.ir/article_513211_575e623abb77f8188092c80a3052a23f.pdf
- Shoab R.M., Abdel-Razik A.B., Ibrahim M.M., Al-Kordy M.A., Taha E.H. 2022. Impact of engineered nano silver on plant parasitic nematode and measurement of DNA damage. *Egyptian Journal of Chemistry* 65: 43–51. DOI: 10.21608/ejchem.2021.99035.4605.
- Shokri S., Babaei A.R., Ahmadian M., Arab M.M., Heshami S. 2015. The effects of different concentrations of nano-silver on elimination of bacterial contaminations and phenolic exudation of rose (*Rosa hybrida* L.) in vitro culture. *Acta Horticulturae* 1083: 391–396. DOI: 10.17660/actahortic.2015.1083.49.
- Shuval H., Yarom R., Shenman R. 2009. An innovative method for the control of Legionella infections in the hospital hot water systems with a stabilized hydrogen peroxide-silver formulation. *International Journal of Infection Control* 5: v5:i1. DOI: 10.3396/ijic.V5i1.006.09.
- Skutnik E., Rabiza-Świder J., Jędrzejuk A., Łukaszewska A. 2020a. The effect of long-term cold storage and preservatives on senescence of cut herbaceous peony flowers. *Agronomy* 10; 1631; 14 p. DOI: 10.3390/agronomy10111631.
- Skutnik E., Jędrzejuk A., Rabiza-Świder J., Rochala-Wojciechowska J., Latkowska M., Łukaszewska A. 2020b. Nanosilver as a novel biocide for control of senescence in garden cosmos. *Scientific Reports* 10; 10274; 9 p. DOI: 10.1038/s41598-020-67098-z.
- Skutnik E., Łukaszewska A., Rabiza-Świder J. 2021. Effects of postharvest treatments with nanosilver on senescence of cut lisianthus (*Eustoma grandiflorum* (Raf.) Shinn.) flowers. *Agronomy* 11:215. DOI: 10.3390/agronomy110202215.
- Smilanick J.L., Mansour M., Sorenson D. 2014. Performance of fogged disinfectants to inactivate conidia of *Penicillium digitatum* within citrus degreening rooms. *Postharvest Biology and Technology* 91: 134–140. DOI: 10.1016/j.postharvbio.2013.12.020.
- Sondi I., Salopek-Sondi B. 2004. Silver nanoparticles as antimicrobial agent: a case study on *E. coli* as a model for Gram-negative bacteria. *Journal of Colloid and Interface Science* 275: 177–182. DOI: 10.1016/j.cis.2004.02.012.
- Stewart-Wade S.M. 2011. Plant pathogens in recycled irrigation water in commercial plant nurseries and greenhouses: their detection and management. *Irrigation Science* 29: 267–297. DOI: 10.1007/s00271-011-0285-1.
- Ślesak I., Libik M., Karpińska B., Karpiński S., Miszański Z. 2007. The role of hydrogen peroxide in regulation of plant metabolism and cellular signalling in response to environmental stresses. *Acta Biochimica Polonica* 54: 39–50. DOI: 10.18388/abp.2007.3267.
- Szpunar-Krok E., Jańczak-Pieniążek M., Skrobacz K., Bobrecka-Jamro D., Balawejder M. 2020. Response of potato (*Solanum tuberosum* L. plants to spraying by hydrogen peroxide. *Sustainability* 12: 2469. DOI: 10.3390/su12062469.
- Thakulla D., Dunn B.L., Goad C., Hu B. 2022. Timing and rates of two products using hydrogen peroxide (H₂O₂) to control algae in Ebb and Flow hydroponic system. *HortScience* 57(1): 32–39. DOI: 10.21273/hortsci16193-21.
- Thakur M., Chandel A., Guleria S., Verma V., Kumar R., Singh G. et al. 2022. Synergistic effect of graphene oxide and silver nanoparticles as biostimulant improves the postharvest life of cut flower bird of paradise (*Strelitzia reginae* L.). *Frontiers in Plant Science* 13: 1006168. DOI: 10.3389/fpls.2022.1006168.
- Torres R., Valentines M.C., Usall J., Viñas I., Larrigaudiere Ch. 2003. Possible involvement of hydrogen peroxide in the development of resistance mechanisms in ‘Golden Delicious’ apple fruits. *Postharvest Biology and Technology* 27: 235–242. DOI: 10.1016/S0925-5214(02)00110-2.
- Tung H.T., Bao H.G., Buu N.Q., Chau N.H., Nhut D.T. 2022. The use of silver nanoparticles as a disinfectant and media additive in plant micropropagation. In: Nhut D.T., Tung H.T., Yeung E.C.-T. (Eds.), *Plant tissue culture: new techniques and application in horticultural species of tropical region*. Springer, pp. 287–302. DOI: 10.1007/978-981-16-6498-4_14.

- Türküsay H., Tosun N. 2005. The effects of hydrogen peroxide treatments on tomato cancer disease (*Clavibacter michiganensis* ssp. *michiganensis* (Smith) Davis et al). Ege Üniversitesi Ziraat Fakültesi Dergisi 42(2): 45–56. [in Turkish with English abstract]
- Uchida A., Takabe T., Takabe T., Jagendorf A.T. 2006. Induction of biosynthesis of osmoprotectants in higher plants by hydrogen peroxide and its application to agriculture. In: Rai A.K., Takabe T. (Eds.), Abiotic stress tolerance in plants. Springer, pp. 153–159.
- Ukuku D.O. 2004. Effect of hydrogen peroxide treatment on microbial quality and appearance of whole and fresh-cut melons contaminated with *Salmonella* spp. International Journal of Food Microbiology 95: 137–146. DOI: 10.1016/j.ijfoodmicro.2004.01.021.
- Webber C. III, Sandtner S., Webber C. 2007. Hydrogen peroxide as a soil amendment for greenhouse nasturtium production (*Trapaeolum majus* L.). Proceedings of Horticultural Industry Show 26: 140–144.
- Wilson R. 2011. Continuous dosing of Huwa-San DW to domestic water system in the United Kingdom. <https://www.koetsplumbing.com/uploads/files/HSP%20DW%20Continuous%20Dosing%20-%20Wilson.pdf>
- Włodarek A., Badełek E., Robak J. 2015. The influence of various products on storage potential of Chinese cabbage (*Brassica pekinensis*) Rupr. Zeszyty Problemowe Postępów Nauk Rolniczych 583: 125–132. [in Polish with English abstract]
- Włodarek A., Sobolewski J., Robak J. 2016. Possibilities of integrated protection against *Alternaria* leaf spot (*Alternaria* ssp.) of Chinese cabbage using three different groups of plant protection products. Progress in Plant Protection 56: 150–154. DOI: 10.14199/ppp-2016-025.
- Wojdyła A. 2011. The possibility of using the preparation Huwa-San TR50 in the protection of ornamental plants against leaf pathogens. Progress in Plant Protection 52: 106–111. [in Polish with English abstract]
- Wojdyła A. 2016a. Możliwość wykorzystania naturalnych i syntetycznych produktów w ochronie róży przed *Podosphaera pannosa* (Wollr.) de Bary. Zeszyty Problemowe Postępów Nauk Rolniczych 586: 89–98. [in Polish]
- Wojdyła A.T. 2016b. The possibility of using natural and synthetic products in control of the willow rust (*Melampsora epitea* Thüm.) occurring on pussy willow (*Salix caprea*). Progress in Plant Protection 56: 236–240. DOI: 10.14199/ppp-2016-040. [in Polish with English abstract]
- Wojdyła A.T., Nowak J.S., Bocianowski J., Wiśniewski J., Waszkiewicz E. 2022. Effect of hyacinth treatment by hydrogen peroxide stabilized with silver and some fungicides on the fungal infection of substrate and bulbs and on plant growth and development. Agronomy 12: 2894. DOI: 10.3390/agronomy12112894.
- Wu C., Shortt B.J., Lawrence E.B., Leon J., Fitzsimmons K.C., Levine E.B. et al. 1997. Activation of host defense mechanisms by elevated production H₂O₂ in transgenic plants. Plant Physiology 115: 427–435. DOI: 10.1104/pp.115.2.427.
- Vanlommel W., Moerkens R., Bosmans L., Lievens B., Rieders H., Wittemans L. et al. 2020. Management of hairy root disease in protected tomato crops: a biological and chemical approach. Acta Horticulturae 1269: 41–50. DOI: 10.17660/actahortic.2020.1269.6.
- Vänninen I., Koskula H. 1998. Effect of hydrogen peroxide on algal growth, cucumber seedlings and the reproduction of shore flies (*Scatella stagnalis*) in rockwool. Crop Protection 17(6): 547–553. DOI: 10.1016/s0261-2194(98)00060-x.
- Yamanaka M., Hara K., Kudo J. 2005. Bactericidal actions of a silver ion solution on *Escherichia coli*, studied by Energy-filtering transmission electron microscopy and proteomic analysis. Applied and Environmental Microbiology 71: 7589–7593. DOI: 10.1007/s00709-018.
- Yan A., Chen Z. 2019. Impacts of silver nanoparticles on plants: a focus on the phytotoxicity and underlying mechanism. International Journal of Molecular Sciences 20: 1003. DOI: 10.3390/ijms20051003.
- Zhao D., Cheng M., Tang W., Liu D., Zhou S., Meng J., Tao J. 2018. Nano-silver modifies the vase life of cut herbaceous peony (*Peonia lactiflora* Pall.) flowers. Protoplasma 255: 1001–1013. DOI: 10.1007/s00709-018-1209-1.
- Zia M., Yaqoob K., Mannan A., Nisa S., Raza G., Rehman R.U. 2020. Regeneration response of carnation cultivars in response of silver nanoparticles under in vitro conditions. Vegetos 33: 11–20. DOI: 10.1007/s42535-019-00074-9.
- Żegliński J. 2006. Stabilizacja nadtlenu wodoru w kserozelu krzemionkowym – badanie oddziaływań składników kompozytu i jego charakterystyka. PhD. Thesis, Akademia Medyczna, Gdańsk. [in Polish]